Single-nucleus proteomics identifies regulators of protein transport

Jason Derks,^{1,2,\overline{\noise}} Tobias Jonson,¹ Andrew Leduc,¹ Saad Khan,¹ Luke Khoury,¹} Mahmoud-Reza Rafiee,¹ Nikolai Slavov^{1,2,⊠}

¹Departments of Bioengineering, Biology, Chemistry and Chemical Biology, Single Cell Center, Northeastern University, Boston, MA 02115, USA

²Parallel Squared Technology Institute, Watertown, MA 02472, USA

 \in Data, code & protocols: scp.slavovlab.net

11

14

The physiological response of a cell to stimulation depends on its proteome configuration. 1 Therefore, the abundance variation of regulatory proteins across unstimulated single cells 2 can be associatively linked with their response to stimulation. Here we developed an ap-3 proach that leverages this association across individual cells and nuclei to systematically Δ identify potential regulators of biological processes, followed by targeted validation. Specifi-5 cally, we applied this approach to identify regulators of nucleocytoplasmic protein transport 6 in macrophages stimulated with lipopolysaccharide (LPS). To this end, we quantified the proteomes of 3,412 individual nuclei, sampling the dynamic response to LPS treatment, and 8 linking functional variability to proteomic variability. Minutes after the stimulation, the pro-9 tein transport in individual nuclei correlated strongly with the abundance of known protein 10 transport regulators, thus revealing the impact of natural protein variability on functional cellular response. We found that simple biophysical constraints, such as the quantity of nu-12 clear pores, partially explain the variability in LPS-induced nucleocytoplasmic transport. 13 Among the many proteins newly identified to be associated with the response, we selected 16 for targeted validation by knockdown. The knockdown phenotypes confirmed the inferences 15 derived from natural protein and functional variation of single nuclei, thus demonstrating 16 the potential of (sub-)single-cell proteomics to infer functional regulation. We expect this 17 approach to generalize to broad applications and enhance the functional interpretability of 18 single-cell omics data. 19

20 Introduction

Single-cell omics methods have rapidly scaled up^{1,2} and enabled the investigation of cellular het-21 erogeneity and the creation of cell atlases³. However, the functional interpretation of such omics 22 data has lagged behind^{4–7}. To investigate the link between proteomic and functional variability 23 across single-cells, we sought to develop an approach that associates the pre-existing differences 24 in proteome configurations to their correspondingly variable cellular responses; such an approach 25 may enable the inference of novel regulatory associations⁸. In particular, we aimed to investi-26 gate how pre-existing proteomic variability influences, and therefore explains the variability of, 27 LPS-induced transport of proteins to and from the nucleus. 28

To quantify this variation between individual cells and individual nuclei, we built upon the con-29 ceptual and technological advances in single-cell mass spectrometry proteomics^{9–21}. Specifically, 30 we used the framework of multiplexed data-independent acquisition (plexDIA) to implement the 31 isotopologous carrier suggestion^{22,23}, which generalizes the concept of isobaric carrier^{24,25} to plex-32 DIA. Isotopologous carriers have already shown promise in some applications^{18,26,27}, and here we 33 used them to enable the first proteomic analysis of individual organelles isolated from human cells. 34 These methodological advances enabled global exploration of protein transport within individual 35 cells, which for decades has been studied by imaging fluorescent proteins²⁸⁻³². Our approach 36 allows for more comprehensive initial discovery, which can expand previous observations that 37 pro-inflammatory stimulation leads to heterogeneous nuclear import of transcription factors^{33,34}. 38 These discoveries can later be examined with much higher temporal resolution using fluorescent 39 imaging^{35,36}. 40

Here, we demonstrate a generalizable approach that enables the inference of functional regula tors from (sub-)single-cell proteomics data. Specifically, we identified proteins regulating subcel lular transport, including the differential contributions of the subunits of the nuclear pore complex.
 Our inferences were derived from natural protein variation across single cells and nuclei. Subse quent validation of these results by targeted knockdown experiments provide direct evidence for
 the functional relevance of our inferential approach.

47 **Results**

48 **Protein transport in bulk macrophage populations**

To first study protein transport between the nucleus and the rest of the cell in bulk populations, six biological replicates were generated using a model system of macrophage-like cells derived from THP-1 monocytes treated with phorbol 12-myristate 13-acetate (PMA). Their nuclei were extracted with mild detergent using methods adapted from an established workflow³⁷. This physical isolation produced sufficient nuclear enrichment as demonstrated by a 5 - 25-fold depletion of proteins from non-nuclear cellular compartments from the nuclei, Extended Data Fig. 1a.

Having validated the enrichment of the isolated nuclear fractions, we next sought to quantify 55 the LPS-induced dynamics of protein transport. To this end, we quantified differential protein 56 abundances for nuclei and whole-cells using MS-EmpiRe³⁸. The abundance changes in the whole 57 cells reflect protein synthesis, degradation, and secretion; changes at the nuclear level also include 58 nucleocytoplasmic transport. As expected, changes mediated by protein synthesis and degradation 59 are slow compared to the kinetics of nucleocytoplasmic transport. Indeed, 60 minutes after LPS-60 stimulation, the changes are dominated by transport with 15.7% of nuclear proteome exhibiting 61 differential abundance compared to only 2.3% of the whole-cell proteome, Fig. 1a. This difference 62 is even more pronounced for earlier time points where changes affect less than 0.1% of the whole-63 cell proteome 30 minutes after LPS stimulation, but 9.3% of the nuclear proteome. 64

Given that hundreds of proteins without innate immunity associations significantly changed 65 their nuclear abundance in response to LPS, we aimed to evaluate the temporal continuity of each 66 protein to further assess the confidence in these findings. As a quantifiable continuity metric, 67 we calculated a ranked version of von Neumann's ratio (RVN)³⁹ for each protein as shown in 68 Extended Data Fig. 2a. Indeed, proteins found to be differentially abundant at 5% FDR for nuclei 69 and whole-cells were mostly monotonic and generally continuous, beyond the monotonicity of a 70 simulated null model where the same number of proteins were assigned random fold-changes and 71 RVN ratios were calculated, Extended Data Fig. 2b,c. Therefore, the large proportion ($\sim 16\%$) of 72 nuclear proteins changing significantly in response to LPS exhibit mostly continuous dynamics, 73 which supports the validity of the findings. 74



Figure 1 | Protein dynamics in whole cells and nuclei reveal coordinated spatiotemporal control of biological processes in bulk populations of macrophage-like cells. a Percent of differentially abundant proteins (5% FDR) in nuclear (grey) and whole-cell (black) proteomes after 10, 30, or 60 minute LPS treatments. b Differentially abundant proteins in the nucleus were analyzed to investigate the relationship between protein mass and transport kinetics. Transport dynamics for each protein were interpolated across the four time points, and ordered with respect to protein mass. The heatmap is colored according to each protein's absolute-valued magnitude of protein transport. Using a moving median approach to average the inherent biological variability, the data suggest smaller proteins achieve half of their total transport at earlier time points than larger proteins. The dashed black line marks a commonly recognized theoretical limit of passive diffusion (40 kDa). \mathbf{c} Time-series data over the course of the LPS-treatment for differentially abundant proteins in the nucleus; pro-inflammatory transcription factors are labeled and highlighted in red, and other proteins with >4-fold change are highlighted in blue. d Protein set enrichment analysis of nuclei and whole-cell bulk samples for NT, 10 minute, 30 minute, and 60 minute LPS treated samples. Brackets correspond to the number of proteins included in computing the enrichment of Gene Ontology terms. e Gene Ontology terms associated with mediating gene-expression, which significantly change in nuclear abundance in response to LPS, are plotted. The 95% confidence intervals are derived from the change of all proteins which correspond to the respective Gene Ontology term. f Proteins were grouped thematically and plotted to show their change in nuclear abundance; all proteins were differentially abundant at 5% FDR in at least one time point. g Nucleoporins which were differentially abundant in the nucleus in at least one time point are plotted to display their change in nuclear and whole-cell abundance; TPR is highlighted in red.

75 Mass dependence of nucleocytoplasmic transport

The kinetics of nucleocytoplasmic transport is mass-dependent, and we sought to investigate this 76 dependence in our data. There are two mechanisms by which transport occurs: passive diffusion 77 and active transport^{40,41}. Molecules with masses less than 40 kDa have been reported to passively 78 diffuse through nuclear pores with kinetics that are mass-dependent⁴². Kinetics of active transport 79 are also dependent on the size of the cargo, but to a lesser degree, as the energy barrier is reduced 80 through interactions with chaperones^{40,43}. We investigated this relationship in our data and found 81 that protein transport was indeed negatively correlated with protein mass, as shown in Extended 82 Data Fig. 3. To increase the time resolution of this analysis, we interpolated protein transport 83 across the 4 time points and found that smaller proteins achieved half their total transport at an 84 earlier time than larger proteins, as shown as a heatmap and scatter plot in Fig. 1b. While this anal-85 ysis cannot conclude whether these trends reflect passive diffusion or simply mass-dependence in 86 active transport, the results are consistent with more recent findings of passive diffusion occur-87 ring without a precise size threshold⁴⁴. Irrespective of the mechanism, these data corroborate the 88 mass-dependence of protein transport kinetics globally in the natural response of macrophages to 89 LPS. 90

⁹¹ Spatiotemporal control of inflammatory response in bulk cell populations

While pro-inflammatory transcription factors are known to be imported to the nucleus in response 92 to LPS, we identify hundreds of additional proteins significantly changing in nuclear abundance, 93 many of which have no prior association with LPS-response. As expected, transcription factors 94 NF- κ B1, REL, RELA, and FOS increase in nuclear abundance by approximately 4-fold within 95 10-60 minutes of LPS-treatment^{29,32,33,45}, as shown in Fig. 1c. Interestingly, hundreds of additional 96 proteins experience monotonic or generally continuous changes in nuclear abundance; one of these 97 proteins, PAXX—a protein required for Non-Homologous End-Joining (NHEJ) DNA repair^{46,47}, 98 reached a 40-fold increase within 60 minutes of LPS-stimulation. This protein acts as a scaffold 99 for the accumulation of XRCC5 and XRCC6 to facilitate NHEJ DNA repair⁴⁸. Likewise, XRCC5 100 and XRCC6 also increased significantly in the nucleus by 60 minutes, Extended Data Fig. 4. 101

To characterize the effects of protein synthesis, degradation, and secretion, we computed pro-102 tein set enrichment on whole-cell proteomes; similar analysis of nuclear proteomes would ad-103 ditionally reflect nucleocytoplasmic transport, Fig. 1d. LPS-stimulation induced whole-cell pro-104 teome enrichment of Gene Ontology (GO) terms associated with gene-expression, such as "trans-105 lation," "spliceosomal complex," and "ribosome biogenesis." Interestingly, changes to the nuclear 106 proteome revealed spatiotemporal changes consistent with this whole-cell enrichment; specifically 107 GO terms whose sites of action are in the nucleus, such as "general transcription initiation fac-108 tor activity" and "ribosome biogenesis" increased in nuclear abundance, while GO terms whose 109 sites of action are outside of the nucleus, such as "cytosolic ribosome" decreased in nuclear abun-110 dance. These changes which reflect coordinated spatiotemporal rearragment of proteins involved 111 in mediating gene-expression are shown for five GO terms in Fig. 1e. Together, the whole-cell and 112 nuclear proteomic data provide complementary evidence which suggests macrophages increase 113 their capacity to mediate gene-expression in response to LPS, as observed through increases in the 114 absolute abundances of and the spatiotemporal rearrangement of proteins involved in this process. 115

To more closely investigate the concordance of changes in nuclear biological processes, dif-116 ferentially abundant proteins were grouped thematically by function and plotted in Fig. 1f. We 117 found concerted dynamics for proteasomal subunits and proteins associated with stress granule as-118 sembly, which decreased in nuclear abundance in response to LPS. Proteins associated with DNA 119 repair, RNA processing, and nucleoporins were generally found to increase in nuclear abundance 120 following LPS stimulation. Intriguingly, of all nucleoporins which changed significantly in nuclear 121 abundance, only TPR—a negative regulator of NPC assembly⁴⁹—decreased, Fig. 1g. Whole-cell 122 abundances of the same nucleoporins were generally found to increase, including TPR. Taken 123 together, these findings are consistent with LPS-induced upregulation of NPC assembly. 124

125

Benchmarking protein quantification of single nuclei

¹²⁶ Using natural variation across single-cells to identify regulators of protein transport requires accu-¹²⁷ rate quantification of many proteins in single nuclei. To accomplish this, we used highly parallel ¹²⁸ sample preparation by nPOP^{19,50} and the plexDIA framework combined with an isotopologous ¹²⁹ carrier. This methodology is analogous to isobaric SCoPE-MS^{15,24,51}, thus we term it SCoPE-

DIA (Single Cell ProtEomics by Data-Independent Acquisition). As we previously suggested and 130 demonstrated with plexDIA, this framework uses the chromatographic coelution of peptides from 131 different samples to increase data-completeness^{22,23}. Thus, protein identification across all samples 132 can be enhanced by including a highly abundant sample, e.g., an isotopologous carrier, in parallel 133 to single-cells or single-organelles. This benefit was evident with the introduction of $plexDIA^{22}$ 134 and has since been reproduced for single-cell analysis¹⁸. However, complex isotopologous carriers 135 might increase interferences and thus decrease quantitative accuracy. To evaluate these trade-offs 136 and find an optimal carrier level for acquiring single-nucleus data, we created and acquired data 137 from a mixed-species spike-in of *H. sapiens* nuclei and *S. cerevisiae* run with 0x, 1x, 5x, 10x, 25x, 138 or 50x carrier amounts, as shown in Extended Data Fig. 5a. We found that intersected protein-level 139 quantitation was comparable across all carrier-levels, Extended Data Fig. 5b. Thus, the potential 140 for interference by larger carriers has little effect on quantitative accuracy in our experiments, 141 which is likely due to the lower proteomic complexity of nuclei as compared to whole cells. 142

To improve quantitative accuracy, the carrier channel was used as a reference to estimate quan-143 titative compression and remove poorly quantified precursors similar to previous filters¹⁵. We 144 applied this filter at various levels to converge upon an optimal balance between coverage and ac-145 curacy, Extended Data Fig. 5c-e. Using this strategy, we compared the coverage and quantitative 146 accuracy and for all proteins identified at various carrier levels and found reduced quantitative ac-147 curacy at higher carrier levels, Extended Data Fig. 5f. This likely reflects the increasing proportion 148 of lowly abundant proteins, which naturally suffer from noisier quantitation. Indeed, the quantita-149 tive accuracy for the subset of proteins quantified across all carrier levels is high even for the 50x 150 carrier, Extended Data Fig. 5b. To summarize, large nuclear carriers were found to not worsen 151 quantitative accuracy, but rather enable quantification of additional lowly abundant proteins that 152 are naturally less well-quantified. Therefore, we chose to acquire single-nucleus data with 25x and 153 50x nuclear proteome carriers. 154

155

Variability in LPS-induced nucleocytoplasmic transport across single nuclei

Next, we sought to infer potential regulators of nucleocytoplasmic transport from the natural vari ability of macrophage proteomes prior to the LPS stimulation and its regulatory impact on the LPS

response. Using SCoPE-DIA, we acquired a dataset consisting of 3,412 nuclei across four biological replicates, with a median of 1,366 proteins quantified per nucleus, Fig. 2a. After filtering to remove nuclei and precursors with poor quantitation (Extended Data Fig. 6a-b), and nuclei with insufficient nuclear enrichment (Extended Data Fig. 6c-d), we retained 2,997 nuclei with a median of 1,287 proteins per nucleus to be used in downstream analyses, Fig. 2a.

Projecting the single-nuclei proteomes onto their principal components produced a partial sep-163 aration between NT and LPS-treated populations, Fig. 2b. Aside from measurement error, three 164 compounding sources of variability may result in this observed partial separation: pre-existing 165 variability of protein abundances at the whole-cell level, pre-existing variability in subcellular pro-166 tein localization, and variability in nucleocytoplasmic transport in response to LPS. We sought to 167 overcome the first two sources of variability and investigate the latter, heterogeneity of nucleocyto-168 plasmic transport. This presents a challenge as the pre-existing protein variability is large relative 169 to the magnitude of protein transport induced by a short LPS exposure. 170

To explore the variability of LPS-induced nucleocytoplasmic transport, we derived a metric 171 aiming to quantify the amount of protein transport experienced by each nucleus; however, given 172 that we did not measure proteomes of the same nuclei before and after stimulation, our estimate 173 cannot directly quantify protein transport. As a substitute, we derived a 'transport score' metric, 174 which quantifies the deviation of single LPS-treated nuclear proteomes from the proteome distri-175 bution of untreated single nuclei. To mitigate noise, we used our bulk data to differentially apply 176 weights based on transporting proteins, as illustrated in Extended Data Fig. 7. As expected, the 177 resulting distributions of transport scores for single nuclei indicate increased nucleocytoplasmic 178 transport for longer time-periods of LPS treatment, Fig. 2c. 179

To link proteomic variability to functional variability in protein transport, we computed correlations of relative protein abundances to transport scores for all LPS-treated single nuclei, Fig. 2d; two proteins, NUP205 ($\rho = 0.28$) and VIM ($\rho = -0.63$), which are among the most (anti-)correlated to protein transport, are highlighted. Protein-set-enrichment results on this ordered vector indicate associations to cell division, cell adhesion, and other processes shown in Extended Data Fig. 8. To explore whether these associations are driven by pre-existing proteomic variability or LPS-induced changes, we compared the protein abundance distributions before and after



Figure 2 | Associating single-nucleus proteomic variability to heterogeneous LPS-induced nucleocytoplasmic protein transport. a LC-MS/MS single-nucleus proteomics data was acquired with SCoPE-DIA for 3412 nuclei, quantifying a median of 1366 proteins per nucleus. Single nuclei which were insufficiently prepared, acquired, or depleted of non-nuclear proteins were removed, yielding 2997 nuclei with a median of 1287 proteins per nucleus post-filtering; these nuclei were used for downstream analyses. **b** Weighted PCA produced partial separation between LPS-treated (10, 30, and 60 minute) nuclei and not-treated (NT) nuclei. **c** Distribution of transport scores from single nuclei. Macrophages treated with LPS for longer durations yielded nuclei which had undergone more protein transport. **d** Rank sorted correlations between the transport score of each nucleus and its proteome. The underlying data used to compute these associations is shown for two proteins, NUP205 and VIM. **e** Distributions of relative protein abundances of NUP205 and VIM in the NT and LPS-treated populations. High values suggest pre-existing variability is much larger than LPS-induced changes. **f** Distribution of overlaps for all proteins between NT and LPS-treated populations of single nuclei.

LPS treatment, Fig. 2e. The results indicate that LPS-induced changes are small relative to the natural variation across untreated nuclei, Fig. 2e,f. Therefore, our associations are dominated by the pre-exisiting variation across proteomic configurations; this initial variability likely influences the amount LPS-induced protein transport each cell undergoes. Thus, associations between singlenucleus protein abundances and transport scores may be used to identify proteins whose abundance affected nucleocytoplasmic protein transport.

¹⁹³ More nuclear pore complexes, more transport

Nearly all nucleoporins were positively correlated to nucleocytoplasmic protein transport, Fig. 3a, 194 suggesting nuclei with more nuclear pore complexes (NPCs) experience more transport. The only 195 exception was TPR, which has been reported to negatively regulate NPC assembly, as validated 196 through knockdowns of TPR yielding cells with more assembled NPCs⁴⁹, as well as in patient 197 fibroblasts with reduced TPR abundances⁵². Interestingly, our data support these previous results, 198 as TPR is the only nucleoporin whose abundance was negatively associated with protein trans-199 port. To more clearly investigate this relationship, we collapsed all nucleoporin abundances to the 200 complex-level, and correlated these NPC abundances to transport scores in LPS-treated nuclei (n 201 = 2,237) and found a relatively strong and highly significant association ($\rho = 0.48, P < 10^{-15}$), as 202 shown in Fig. 3b. Indeed, this suggests a highly significant association between NPC abundance 203 and nucleocytoplasmic transport. 204

Having uncovered this association between transport and NPC abundance in single nuclei, we 205 sought to additionally investigate whether the mass-dependence of nucleocytoplasmic transport 206 also changes as the number of NPCs varies across single nuclei; specifically, the nuclear envelope 207 should become more permeable to passive diffusion at higher pore densities. To test this hypoth-208 esis, we first quantified the mass dependence of protein transport for each nucleus by regressing 209 the molecular masses of proteins on their absolute deviations from the NT-population of single 210 nuclei-a substitute for measuring protein-specific transport. The slopes of these regressions were 211 plotted against corresponding NPC abundances in single nuclei, Fig. 3c. The results revealed a 212 modest association across all 2,237 LPS-treated nuclei ($\rho = -0.17$) with high statistical signifi-213 cance ($P < 10^{-14}$). This association indicates that nucleocytoplasmic transport of smaller proteins 214



Figure 3 | Nucleoporin abundances and their associations to LPS-induced nucleocytoplasmic transport. a Nucleoporins are variably associated with LPS-induced nucleocytoplasmic protein transport. b Single nuclei with more nuclear pore complexes experience more LPS-induced nucleocytoplasmic protein transport ($P < 10^{-15}$, $\rho = 0.48$). c For each single LPS-treated nucleus, a slope of the absolute value of differentially abundant proteins vs protein mass was computed to quantify mass-dependence of transport for each single nucleus; single nuclei which had the least and greatest abundance of nuclear pore complexes are shown at the top of the panel. The resulting slopes were plotted against their respective nuclear pore complex abundances, as shown at the bottom of the panel. The data suggest protein transport becomes increasingly mass-dependent, favoring transport of smaller proteins, when nuclear pore complex abundances are high ($P < 10^{-14}$, $\rho = -0.17$). d Predicted human nuclear pore complex from Mosalaganti *et al.*, with nucleoporins colored by their correlations to transport score. e Distribution of each nucleoporin's correlation to transport score grouped by whether it is a peripheral or scaffold protein in the NPC (P = 0.037). g Nucleoporins' correlations to transport are associated with their respective half lifes (half-life data from Mathieson *et al.*).

increasingly outpaced transport of larger proteins, at least in part as a function of NPC abundance,
 Fig. 3c. Given that passive diffusion is highly mass-dependent, it may suggest that single nuclei with more NPCs experience more passive diffusion, thus providing empirical support for this
 theoretical expectation.

While the abundances of nearly all nucleoporin proteins correlate positively with transport 219 scores, the magnitude of this correlation varies across nucleoporin proteins, as shown in Fig. 3a. 220 To explore this variability, we investigated what might be associated with these differences. To 221 visually inspect the structural dependence of the correlations, the predicted structure of the human 222 nuclear pore complex from Mosalaganti et al.⁵³ was colored according to each proteins' correlation 223 to the transport score, Fig. 3d. This was investigated more directly by grouping nucleoporins 224 based on their localizations inside the nuclear pore complex, as shown in Fig. 3e, and according to 225 whether the nucleoporins were peripheral or scaffold proteins of the NPC⁵⁴, as shown in Fig. 3f. 226 The abundances of scaffold proteins were significantly more associated with nucleocytoplasmic 227 transport than peripheral proteins of the NPC (P = 0.037). This variation across nucleoporins is 228 further supported by a significant correlation with protein half-lives derived from Mathieson et al.'s 229 SILAC turnover data⁵⁴ ($\rho = 0.56$, P = 0.0018), as shown in Fig. 3g, as scaffold nucleoporins have 230 been reported to have longer half-lives than peripheral nucleoporins^{54,55}. These results suggest the 231 observed variability in each nucleoporins' correlation to transport is significantly associated with 232 the nucleoporin's localization inside the NPC and their respective half-lives. 233

234

Targeted perturbations validate the single-nucleus derived associations

Having used single-nucleus proteomics data to infer novel regulators of LPS-induced nucleocyto-235 plasmic protein transport, we aimed to directly test these potential enhancers and suppressors of 236 nucleocytoplasmic transport through targeted perturbations. The highest ranking hypotheses from 237 the single-nucleus proteomics analysis include proteins directly involved in transport such as nu-238 cleoporins and ribosomal export proteins such as MDN1⁵⁶. However, many other proteins without 239 functional annotations for protein transport or innate immunity are also highly (anti-)correlated 240 to transport. Perhaps some of these proteins influence LPS-induced protein transport, and these 241 functional associations present discovery opportunities. We sought to test these associations ex-242

perimentally through directed perturbations using siRNA mediated knock downs.

243

To test potential transport regulators, we chose to knock down 16 proteins spanning the con-244 tinuum shown in Fig. 4a and quantified the change in LPS-induced nucleocytoplasmic protein 245 transport. The knockdowns were performed using siRNAs as shown in Fig. 4b, and the de-246 crease of the corresponding gene products was validated in whole-cell and nuclear bulk lysates, as 247 shown in Supplementary Fig. 1 and Supplementary Fig. 2. To quantify how each knockdown af-248 fected LPS-induced nucleocytoplasmic protein transport, slopes were calculated from fold-changes 249 (LPS/NT) between the original bulk data and each siRNA condition, using the negative control 250 (non-targeting) siRNA as a reference for each biological replicate, as shown in Fig. 4b. Results for 251 individual biological replicates are shown in Supplementary Fig. 3. With this strategy, we aimed 252 to quantify the global change in nucleocytoplasmic transport as a result of each knockdown. 253

Associating relative protein abundances to function in single cells may serve as a means of 254 inferring regulatory potential. We sought to evaluate the predictive power of these associations 255 by computing the significance of protein correlations to transport from the single-nuclei data and 256 the significance of the change in transport from corresponding knockdowns, Fig. 4c. Indeed, as 257 a result of the knockdowns, protein transport for 13/16 of the conditions was affected in the ex-258 pected direction. Specifically, knockdowns of potential transport suppressors increased transport 259 in 7/7 cases; knockdowns of potential transport enhancers decreased transport in 6/9 cases. The 260 protein correlations to transport scores from the single-nucleus data were highly predictive of the 261 functional effects of the knockdowns ($\rho = -0.77, P < 5 \times 10^{-4}$). The slopes, which were used 262 for determining the change in LPS-induced nucleocytoplasmic transport, were computed from the 263 100 most differentially abundant proteins from the original bulk data. Various filtering levels were 264 applied to examine the robustness of the findings, as shown in Extended Data Fig. 9, and all three 265 levels showed similar trends. 266

Knockdowns of IMMT, RPSA, HSD17B4, SND1, DHX15, and VIM were found to significantly increase transport, whereas knockdowns of RBM6 and MYBBP1A were found to significantly reduce transport. Therefore, 8/16 knockdowns significantly changed LPS-induced nucleocytoplasmic transport, 7/8 of which changed nucleocytoplasmic transport in the expected direction. The knockdown which most significantly affected transport was IMMT, a mitochondrial protein



Figure 4 | **Genetic perturbations validate single-nucleus-derived protein associations**. **a** Correlations of protein abundances to transport score from single-nucleus data. 16 proteins were selected to be knocked-down and colored with green or purple depending on whether it is associated with increased or decreased transport. **b** Workflow for siRNA-mediated knockdowns in THP-1 monocytes, followed by differentiation to macrophages. A negative control (non-targeting) siRNA was included in each biological replicate, and used to compute a change in transport as the result of each siRNA knockdown (KD); change in transport was calculated as the difference in slopes between the negative control siRNA and a gene-targeting siRNA. **c** Here we evaluate the predictive power of the single-nucleus data were used to compute significance of correlations to transport scores, shown on the y-axis; the x-axis measures the significance of how the knockdown affected nucleocytoplasmic protein transport. The single-nucleus associations were highly predictive of the functional effects in genetic perturbations (*P* = 0.00046, $\rho = -0.77$). **d** Significance of how individual gene knockdowns affect nucleocytoplasmic transport. Knockdowns of IMMT, RPSA, HSD17B4, SND1, DHX15, and VIM were found to globally increase LPS-induced nucleocytoplasmic protein transport; knockdowns of RBM6 and MYBBP1A were found to decrease transport.

without prior association to innate immunity or nucleocytoplasmic protein transport. Whole-cell
 lysates of these knockdowns complementarily indicate a gradient of unpreparedness and primed ness to respond to LPS and facilitate nucleocytoplasmic transport, as shown in Extended Data
 Fig. 10. These findings validate the potential of (sub-)single-cell proteomics to identify novel
 regulators of biological functions.

Discussion

277

Identifying functional regulators of complex biological processes is an ongoing challenge in pro-278 teomics data interpretation. Here we demonstrate an interpretation approach that exploits the natu-279 ral variability inherent in a population of single cells to empower inference of functional regulation. 280 It is conceptually similar to the identification of functional regulators through perturbation-induced 281 variability in bulk samples⁵⁷ but instead utilizes the natural variability across individual cells. In 282 our approach, we assumed that the initial proteomic states of single macrophages would explain 283 the variability in LPS-induced nucleocytoplasmic protein transport. Operating under this principle, 284 we quantified a metric (the transport score) from which we could evaluate the influence of resting 285 proteomic variability on this response. We found that simple biophysical constraints, such as the 286 quantity of nuclear pores, partially explain the variance in nucleocytoplasmic transport. Interest-287 ingly, the abundance of scaffold subunits is more strongly associated with transport rates than the 288 abundance of peripheral subunits, which is consistent with modular and specialised structures of 289 the nuclear pore complex⁵⁸. 290

Beyond the role of nuclear pores, our analysis identified hundreds of additional proteins, many 291 of which without annotated associations to innate immunity or protein transport, yet whose abun-292 dances were strongly correlated to the magnitude of nucleocytoplasmic protein transport expe-293 rienced in single cells. We tested these protein-associations through genetic perturbations and 294 found that the associations derived from single-nucleus data were, as a whole, highly predictive 295 of their respective effects on LPS-induced nucleocytoplasmic transport. This work demonstrates 296 an approach which uses (sub-)single cell proteomics to infer functional regulators of complex bio-297 logical processes, such as LPS-induced nucleocytoplasmic transport. We expect approaches such 298 as this will generalize to broad applications and support the functional interpretation of single-cell 299

³⁰⁰ proteomics data.

301 Methods

302 Cell culture

THP-1 monocytes (TIB-202, ATCC) were cultured in RPMI-1640 Medium (R8758, Sigma-Aldrich) 303 supplemented with 10% fetal bovine serum (10439016, Gibco) and 1% penicillin-streptomycin 304 (15140122, Gibco) and grown at 37°C and 5% CO₂. Cells were transferred to 100x15mm Nunclon 305 dishes (150350, Thermo Scientific) at a density of approximately 200,000 cells/mL and volume of 306 10 mL. Differentiation proceeded following an established protocol by Lund et al.⁵⁹; THP-1 cells 307 were incubated in the presence of 25 nM phorbol 12-myristate 13-acetate (PMA) for 48 hours, then 308 cells were detached with accutase, mixed to prevent any batch-effects resulting from differentia-309 tion occurring in isolated plates, washed 2-times with PBS, centrifuging at 300g for 5 min to pellet 310 for washes. The cells were then allowed to incubate in growth media without PMA for 24 hours, 311 and in this time reattach to fresh 100x15mm Nunclon dishes. The resulting M0 macrophages were 312 either left not-treated (NT), or treated with 1 μ g/mL lipopolysaccharides from *Escherichia coli* 313 O111:B4 (L4391, MilliporeSigma) for 10 minutes, 30 minutes, or 60 minutes. 314

315 **Protein knockdowns**

Proteins of interest were knocked down via reverse transfection of Silencer siRNAs (Ambion) 316 and Silencer Select siRNAs (Ambion) into THP-1 monocytes using methodology adapted from 317 manufacturer instructions for RNAi transfection. Silencer siRNAs were transfected in 12-well 318 Nunclon plates and Silencer Select siRNAs were transfected in individual 60 mm dishes. For 319 Silencer siRNAs, 10 pmol of siRNA was complexed with 3 μL of RNAiMAX. For Silencer Select 320 siRNAs, 55 pmol of siRNA was complexed with 17 μL of RNAiMAX (Invitrogen) in 552 μL of 321 serum-free RPMI (Sigma) in a 60 mm Nunclon dish (Thermo). siRNAs were complexed with the 322 RNAiMAX lipid reagent for 5 minutes at room temperature. THP-1 monocytes were added in 323 RPMI supplemented with 10% fetal bovine serum (FBS) and 1% penicillin-streptomycin to reach 324 a final density of 300,000 cells per mL. Samples were cultured at 37°C and 5% CO₂ for 24 hours 325

before adding PMA, at which point the same aforementioned cell culture process was followed.

327 Nuclear isolation

326

Macrophages were detached from Nunclon dishes with accutase on ice for approximately 20 min-328 utes. The cells were washed twice with 1x PBS to remove media, centrifuging at 300g for 5 329 min each time. The cell concentration was measured with a hemocytometer. 15% of cells were 330 retained as whole-cells for downstream analysis by resuspending the whole-cell pellet in LC-MS-331 grade water to a concentration of 3,000 cells/ μL . These whole-cells were then stored at -80°C. 332 The remaining 85% of cells were resuspended in 1 mL of ice-cold Lysis Buffer (0.1% NP-40, 250 333 mM sucrose, 25 mM KCl, 5 mM MgCl₂, 10 mM HEPES (pH = 7.4)) in a 2 mL Eppendorf tube, 334 and left to lyse on ice for 20 min. At this point in the protocol, the cells are lysed but the nuclei are 335 unlikely to be free from cellular debris and other organelles. To purify the nuclei, the 1 mL nuclei 336 suspension was sheared through a 25G needle with even pressure, avoiding bubbling and foaming 337 as much as possible. The shearing was repeated for a total of 5 times. Nuclei were visually in-338 spected under a microscope with Trypan Blue staining to ensure sufficient shearing of subcellular 339 organelles from nuclei. 340

After nuclei are sufficiently sheared, the suspension was centrifuged at 1000g for 8 min at 4 °C. 341 The nuclear pellet was resuspended in 500 μ L of 250 mM sucrose, 25 mM KCl, 5 mM MgCl₂, 10 342 mM HEPES (pH = 7.4). The mixture was underlayed with 500 μ L of 350 mM sucrose, 25 mM KCl, 343 5 mM MgCl₂, 10 mM HEPES (pH = 7.4) to form a sucrose cushion, which was then centrifuged 344 at 4,000g for 10 min (4 °C). The supernatant was removed, and the nuclear pellet was resuspended 345 in 200 μ L of LC-MS water, gently flicking to mix. 4 μ L of nuclei suspension was mixed with 4 μ L 346 of Trypan Blue, then counted with a hemocytometer to find the final concentration. 150 µL of the 347 bulk nuclei were stored at -80°C. The remaining 50 µL was diluted to a concentration of roughly 348 400 nuclei/ μ L and used for single-nuclei sorting and sample preparation by nPOP^{19,50}. 349

350 Sample preparation for proteomic analysis

351 Bulk samples

Nuclei and whole-cell fractions, which were frozen at -80°C at a concentration of 3,000 cells/ μ L, 352 were heated to 90°C for 10 minutes as part of the mPOP protocol⁶⁰. Trypsin Gold (V5280, 353 Promega) was added to a final concentration of 20 ng/ μ L, in addition to final concentrations of 354 100 mM TEAB and 0.25 U/ μ L benzonase nuclease (E1014, Millipore Sigma). The cell lysates 355 were digested at 37°C for 18 hours in a thermocycler. Samples were labeled with mTRAQ follow-356 ing manufacturer's instructions. In short, the labels, which arrived suspended in isopropyl alcohol 357 at a concentration of 0.05 U/ μ L (where 1 unit labels 100 μ g), were added in a 1:2 proportion of 358 label:peptide where the peptides were concentrated to 500 ng/ μ L in 100 mM TEAB. The labeling 359 reaction was left at room temperature for 2 hours, and then quenched with 0.25% hydroxylamine 360 for 1 hour. 361

362 Creating a mixed species standard to benchmark SCoPE-DIA

³⁶³ Digested protein lysate from THP-1 macrophage nuclei were mixed with digested *S. cerevisiae* ³⁶⁴ protein lysate (V7461, Promega) to generate three samples. The proportions of Sample A ($\Delta 0$):Sample ³⁶⁵ B ($\Delta 4$) was 4:1 for *S. cerevisiae* and 1:1 for *H. sapiens*. The unlabeled versions of these samples ³⁶⁶ were mixed in a 1:1 ratio to generate a carrier sample which was then labeled with mTRAQ $\Delta 8$. ³⁶⁷ Samples A and B were diluted such that the *H. sapiens* amount was 1 nucleus worth of protein ³⁶⁸ (approximately 45 pg). The carrier sample was pooled with Samples A and B at 0x, 1x, 5x, 10x, ³⁶⁹ 25x, and 50x concentrations to generate the SCoPE-DIA benchmarks.

370 Single nuclei

Single nuclei were prepared for proteomic analysis by nPOP as previously described by Leduc *et al.*^{19,50}. In short, single-nuclei were sorted by CellenOne into 8 nL droplets of 100% DMSO on a fluorocarbon-coated glass slide to lyse. The lysed nuclei were digested with Trypsin Gold at a concentration of 120 ng/ μ L and 5 mM HEPES (pH 8.5) and 200 mM TEAB buffer (pH 8.5). Peptides were then chemically labeled with mTRAQ $\Delta 0$ or $\Delta 4$. Single nuclei were pooled into

sets of $\Delta 0$ and $\Delta 4$, and dispensed into individual wells of a 384 well-plate. The pooled samples in the 384 well-plate were then dried in a speed-vacuum then stored at -80°C until use. Before sample loading for LC-MS/MS, a 25x or 50x nuclei carrier labeled with mTRAQ $\Delta 8$ was pooled into each set to create the final $\Delta 0$, $\Delta 4$, and $\Delta 8$ SCoPE-DIA set. Four single-nucleus biological replicates were prepared and acquired, for a total of 3412 single nuclei.

Data acquisition

All single nuclei and bulk samples were resuspended in 0.01% n-Dodecyl β -d-maltoside (DDM)⁶¹ 382 with 0.1% formic acid. Sample pickup occurred out of a 384 well-plate for single-nucleus sam-383 ples, or from glass vials for bulk samples. Peptides were separated for LC-MS acquistion using a 384 Neo Vanquish UHPLC and 25 cm \times 75 μm IonOpticks Aurora Series UHPLC columns (AUR2-385 25075C18A) at a flow-rate of 200 nL/min. LC was run with the following settings: Direct Injec-386 tion, nano/cap flow, maximum pressure = 1500 bar, maximum pressure change = 1000 bar/min. 387 Sample loading had the following settings: $1\mu L$ injection with 1.2 μL loading volume, "Pressure 388 Control" mode with maximum pressure set to 1450 bar, and fast loading enabled. Wash and equi-389 libration settings: "Pressure Control" mode with maximum pressure set to 1450 bar, equilibration 390 factor = 4.0, and fast equilibration was enabled. 391

The LC gradient proceeded as follows: Ramp from 2.5%B to 6.5%B over 0.2 minutes, to 11.5%B over 0.9 minutes, to 21.0%B over 3.1 minutes, to 31.5%B over 6.2 minutes, to 40%B over 2.8 minutes, to 55%B over 1.7 minutes, to 95%B over 0.65 minutes and then hold at 95%B for 4 minutes. This method allows for approximately 15 minutes of active chromatography, at a throughput of approximately 45 runs/day (90 single nuclei/day with SCoPE-DIA or 135 bulk samples/day with plexDIA) after accounting for sample loading and column equilibration overheads.

All data were acquired on a Bruker timsTOF SCP using captive spray, dia-PASEF scan mode^{62,63}, and positive polarity. The duty cycle consisted of 8 total PASEF frames, 26 Th MS2 windows with 1 Th overlaps. To improve temporal sampling of the MS1-elution profiles which are used for quantification, an MS1 scan was taken every 2 PASEF frames, for a total of 4 MS1 scans per duty cycle. MS2 scan range: 300 m/z-1000 m/z, MS1 scan range: 100 m/z-1700 m/z, 1/K0 start: 0.64, 1/K0 end: 1.20, ramp and accumulation times: 100 ms. The estimated duty cycle time is 1.28 seconds.

Collision energy settings were 20 eV at 1/K0 = 0.60 and 59 eV at 1/K0 = 1.60. Collision RF was set to 2000 Vpp.

406 **Raw data searching with DIA-NN**

Empirical spectral libraries of mTRAQ-labeled nuclei and whole-cells were generated from dia-PASEF runs on the timsTOF SCP, searching with a DIA-NN in-silico-predicited Swiss-Prot *H. sapiens* FASTA (canonical & isoform). Similarly, a combined *H. sapiens* and *S. cerevisiae* library was generated for searching SCoPE-DIA benchmarks.

DIA-NN (version 1.8.1) was used to search raw data as previously implemented^{22,64}. The following search settings were used: {-fixed-mod mTRAQ 140.0949630177, nK}, {-channels mTRAQ, 0, nK, 0:0; mTRAQ, 4, nK, 4.0070994:4.0070994; mTRAQ, 8, nK, 8.0141988132:8.0141988132}, {-original-mods}, {-peak-translation}, {-report-lib-info}, {-qvalue 0.01}, {-mass-acc-ms1 5.0}, {-mass-acc 15.0}, {-mass-acc-quant 5.0}, {-reanalyse}, {-rt-profiling}, and {-peak-height}.

416 **Computational data analysis**

417 **Nuclear enrichment**

To assess nuclear enrichment, the first two bulk biological replicates were acquired in a plexDIA set of whole-cell and nuclear fractions. The fractions were subset for histones, which should only be present in the nucleus, and both fractions were normalized by a scalar such that histone proteins between whole-cells and nuclear fractions were in a 1:1 ratio. Ratios of MaxLFQ protein abundances for protein markers from The Human Protein Atlas for endoplasmic reticulum (ER), nuclear compartments, mitochondria, plasma membrane, cytosol, and Golgi were calculated between nuclear and whole-cell fractions to assess nuclear protein enrichment^{65–67}.

425 Differential protein abundance analysis

426Precursor quantities acquired by plexDIA and analyzed by DIA-NN were corrected for isotopic en-427velope carryover, as previously performed²². These precursor quantities were filtered for Lib.PG.Q. Value428< 0.01 and Q. Value < 0.01 for bulk data from all six biological replicates and their corresponding

nested technical replicates. The bulk data used to generate weights for single-nuclei analyses were
taken from cell-batches that corresponded to those single nuclei, in this case, biological replicates
1, 2, 3, and 6; the bulk data used for single-nucleus weighting were further filtered for Translated
Q-value and Channel Q-value < 0.05.

Differential protein abundance was calculated through the MS-Empire workflow³⁸ for each 433 biological replicate. Precursors used for analysis were required to be quantified in at least two 434 replicates per condition. Data normalization was performed using operations from the MS-Empire 435 workflow, as described by Ammar *et al.*³⁸. Differential abundance analysis collapsed precursors to 436 gene-level annotations; this produced p-values calculated after outlier correction for each protein 437 for each biological replicate; Stouffer's method was applied to collapse multiple p-values from 438 several biological replicates to a single combined p-value. Briefly, this involved converting p-439 values for each protein from each biological replicate to z-scores, multiplying by the sign of the 440 fold-change, summing the signed z-scores, then dividing by the square root of the number of 441 observations for that protein, and finally converting that z-score back to a single p-value for each 442 protein. The resulting p-values were corrected for multiple hypothesis testing using the Benjamini-443 Hochberg (BH) correction⁶⁸. 444

445 **Protein set enrichment analysis**

Protein set enrichment analysis was performed using the g:GOSt R package^{69,70}. An ordered vector 446 of gene names was used as the input. Results were filtered at 1% FDR, and the relative abundance 447 for the Gene Ontology terms was computed across all samples from the intersect of protein abun-448 dances to represent the relative abundance of a given Gene Ontology term. Calculating the relative 449 abundance of a Gene Ontology term from the subset of intersected proteins allowed for fairer com-450 parisons of enrichment between samples. Specifically regarding the analysis of whole-cell bulk 451 knockdowns shown in Extended Data Fig. 10, in some cases this was performed on data in which 452 NT and 60 minute LPS-treated cell-lysates were combined. These samples were experimentally 453 combined for ease of labor and due to the minimal impact of 60 minute LPS-treatment on the 454 whole-cell proteome. 455

456 Evaluating relationship of protein mass and transport

Protein masses were downloaded from Uniprot⁷¹, and matched to proteins which we found significantly increased or decreased in response to LPS (Q-value < 0.05) in nuclear fractions. The absolute value of the log₂ fold-change was plotted with the mass of the protein to statistically evaluate the relationship between protein mass and the amount of transport that occurred as a result of LPS treatment. Because the masses are purely based on gene-encoded sequences, this analysis does not account for changes in mass as the result of proteolytic cleavage or the transport of protein complexes.

Additionally, the dynamics of protein transport were interpolated from the four data points (NT, 464 10 min, 30 min, and 60 min LPS) using a 3rd degree polynomial for each of the proteins that was 465 found to significantly change in at least one of the conditions relative to NT. A total of 300 time 466 points were predicted from the fit, ranging from 0 to 60 minutes. The proteins were arranged from 467 smallest to largest and a moving median of the absolute value of the log₂ fold-change from NT was 468 calculated for each time point from that protein's adjacent 40 smaller and 40 larger proteins. In this 469 way, biological differences between proteins could be averaged to generally assess the influence 470 of protein mass on transport kinetics. For each protein's averaged kinetics, the time when half of 471 the total absolute value log₂ fold-change had occurred was marked. This was plotted against the 472 protein mass and a Spearman correlation was calculated. 473

474 Benchmarking SCoPE-DIA quantification

To benchmark how different levels of carrier affect protein-level quantitation of SCoPE-DIA as shown in Extended Data Fig. 5f, precursors were intersected across all carrier levels, and thus proteins as well; precursor quantities were collapsed to protein-level quantities using MaxLFQ, and then ratio values between $\Delta 0$ and $\Delta 4$ were computed, normalizing *H. sapiens* to a be mediancentered at a 1:1 ratio to account for systematic differences in loading amounts, if any exist. To compare quantitation at different carrier levels for all proteins, as shown in Extended Data Fig. 5f, the same process was performed without intersecting.

482

In regards to compression filtering, acquiring single cell (or nucleus) data with a constant,

known amount of carrier is useful to identify precursors which have systematically interfered
quantification. Precursors which have systematically compressed quantification across all samples
relative to the expected carrier amount, are unlikely to be well-quantified and can be removed. To
identify which precursors to remove, a ratio for each precursor from the single-nucleus channels to
its respective carrier channel was calculated. Precursors with median ratios > 6-fold greater than
the theoretical ratio (e.g. 1:50 is expected for a 50x carrier) were considered to be systematically
poorly quantified and were excluded from downstream analyses.

490 Single-nucleus quality control

To identify which single nuclei were successfully prepared and acquired, precursors were filtered for Channel and Translated Q-values < 0.1 and the number of remaining high quality precursors was counted. Single nuclei with ≥ 50 high quality precursors were retained for downstream analysis as this threshold was sufficiently separable from negative controls. Precursors were required to be quantified in $\geq 1\%$ of single nuclei, and have < 6x ratio compression to be retained for downstream analysis.

Single nuclei were further filtered to ensure sufficient depletion of non-nuclear proteins. To quantify the nuclear purity of each single-nucleus, the nuclear proteomes were scaled to a reference bulk nuclear fraction, and compared to a ratio of whole-cell to nuclear fraction. A 3rd degree polynomial was used to fit a curve for each single-nucleus, and an absolute value of the AUC was computed. This value was used to identify single-nuclei which were insufficiently pure. Only single nuclei with AUCs < 5 were used in downstream analyses.

503 Single-nucleus protein quantification, imputation, and batch correction

⁵⁰⁴ Precursor abundances from single nuclei were normalized to their respective carrier channel, which ⁵⁰⁵ serves as a reference similar to the isobaric SCoPE-MS workflow^{15,24}. For each precursor across ⁵⁰⁶ all carrier samples, the mean abundance was calculated and used to scale the data of the single ⁵⁰⁷ nuclei, to be compatible for MaxLFQ protein quantification⁶⁵. The resulting protein-level data of ⁵⁰⁸ each nucleus was normalized by its respective median protein abundance to account for differences ⁵⁰⁹ in absolute abundances, then each protein was normalized by the mean abundance across single

⁵¹⁰ nuclei. The \log_2 -transformed data was imputed using a weighted kNN approach (k=5), where ⁵¹¹ the weights were proportional to the similarity to its nearest neighbors. Finally, the data was ⁵¹² batch-corrected using ComBat^{72,73} to account for mTRAQ label-biases, LC-batches, and biological ⁵¹³ replicates.

514 Weighted Principal Component Analysis (PCA)

The variance of single-nucleus protein data was weighted based on protein correlations, bulk differential abundance, and bulk continuity of spatiotemporal trends. Effectively these weights assign higher variance based on: 1) how correlated a protein's abundance is to other proteins abundances, 2) how differentially abundant the protein was in the bulk data, and 3) how continuous the trend was in its spatiotemporal response to LPS.

⁵²⁰ Specifically, the bulk differential abundance data computed from MS-Empire³⁸ analysis had ⁵²¹ p-values which where converted to signed Z-scores. Using Stouffer's method they were combined ⁵²² for all the bulk biological replicates that correspond to the single-nucleus replicates, in this case, ⁵²³ biological replicates 1, 2, 3, and 6. The resulting Z-score for each protein was squared, then used ⁵²⁴ to weight the variance of that protein in the single-nucleus data. Additionally, the inverse of the ⁵²⁵ ranked version of the von-Neumman Ratio (RVN) calculated from bulk data was used to weight ⁵²⁶ the variance of each respective protein in the single-nucleus data.

527 **Computing single-nucleus transport scores**

Directly measuring protein transport in a cell using LC-MS/MS was not possible with this experi-528 mental design as we did not measure the same nucleus before and after LPS stimulation. However, 529 we derived a 'transport score' metric to serve as a substitute which serves to quantify the devi-530 ation of single LPS-treated nuclei from the population of single NT nuclei. The transport score 531 accounts for pre-existing variability in the not-treated (NT) population of single nuclei for each 532 protein by converting protein abundances for each LPS-treated nucleus to a Z-score in reference 533 to the NT-population. For each nucleus, the resulting vector of Z-scores is weighted based on 534 the signed-Z-scores of the differential protein abundances in the bulk data. Therefore, a protein's 535 contribution to the transport score is weighted by its significance in the bulk data, where more 536

differential proteins carry greater weights. The mean component of the resulting vector of each nucleus is taken as that nucleus' transport score.

⁵³⁹ Computing protein overlap coefficients between populations of single nuclei

The overlap between the distributions of relative protein abundances between populations of single nuclei was computed to quantify the commonality of protein abundances pre-stimulation and post-LPS stimulation. This overlap coefficient was computed by the overlap function of bayestestR⁷⁴. It is computed by modeling the distribution of relative protein abundances as densities for two populations and then computing the proportion that is shared. For example, a protein that has high-overlap between abundances in NT and LPS-treated populations, is one that is minimally transported relative to its natural variability.

547 Quantifying nuclear pore complex abundances and dependence on mass

To quantify nuclear pore complex abundances in each single-nucleus, relative abundances of nucleoporins were converted to Z-scores and combined for all nucleoporins using Stouffer's method. Mass dependence as a function of NPC abundance was computed for each LPS-treated singlenucleus based on the slopes (protein Z-scores vs log₁₀-transformed protein mass) of differentially abundant proteins in the single-nucleus data.

Investigating the association between nucleoporin half-lives and their correlations to trans port score

⁵⁵⁵ Half-lives were derived from SILAC turnover data of primary monocytes, NK cells, neurons, hep-⁵⁵⁶ atocytes, and B-cells from Mathieson *et al.*⁵⁴. Specifically, the half-lives of all replicates for each ⁵⁵⁷ cell-type was collapsed to the mean, then the mean of all cell-types was collapsed to a final value, ⁵⁵⁸ which was used as the half-life in our analysis.

559 Visualizing the nuclear pore complex

The predicted structure of the human nuclear pore complex (PDB Entry: 7r5k)⁵³ from Mosalaganti *et al.* was used visualize the NPC. The nucleoporins were colored by their respective correlations to transport score using ChimeraX (v1.6.1)⁷⁵.

Quantifying the effect of protein knockdowns on transport

To quantify how each protein knockdown affected LPS-induced nucleocytoplasmic transport, each 564 biological replicate included a matched negative (non-targeting) siRNA control to account for how 565 much transport would have occurred without the knockdown. A slope was calculated from the 100 566 most differentially abundant proteins in the original (no knockdown) bulk data to the data for each 567 knockdown. The fold-changes for proteins used for computing these slopes were weighted by their 568 associated Z-score significance of differential abundances from the original bulk data; this serves to 569 weight each protein's contribution to the slope-calculation based on its significance. The difference 570 in slopes between the negative control (non-targeting) siRNA and the knockdown was computed. 571 This difference in slope is interpreted as the extent to which nucleocytoplasmic transport changed 572 as a result of the knockdown. Additionally, the associated standard error of each slope was used 573 between the negative control and experimental knockdown to derive a Z-score for the significance 574 of the difference in transport. Significance from additional biological replicates were combined 575 using Stouffer's method to arrive at a cumulative Z-score. To assess the significance of the effect 576 on transport of individual knockdowns, this Z-score was converted to a p-value and adjusted for 577 multiple testing hypotheses using a BH-correction. 578

579 Calculating significance of correlation to transport score

To derive a metric for the significance of protein correlations to the transport score, a rank-based normalization approach (Rankit) was used to transform the correlations into a standard normal distribution. The resulting Z-scores were combined using Stouffer's method to arrive at a final Z-score for each protein, as shown on the y-axis of Fig. 4c. This approach was chosen as an alternative to a traditional Z-score operation given that the distribution of correlations to transport

scores were non-Gaussian and have a skewed tail.

586 Availability

585

All data are reported and deposited in accordance with the community guidelines¹³. Raw data, spectral libraries, FASTAs, and DIA-NN search outputs are available at MassIVE: MSV000094829. Code, processed data, supporting files, supplementary figures, and interactive supplementary analyses (e.g. volcano plots and time-series data) are available at scp.slavovlab.net/Derks_et_al_2024.

Acknowledgments: We thank H. Specht, R. Huffman, A. Petelski, D. Perlman, and E. Emmott for 591 discussions pertaining to experimental methodology for single-nucleus proteomics. We thank M. 592 Rout and T. Eeuwen for thoughtful discussions regarding nuclear pore complexes. And finally, we 593 thank I. Zanoni and M. Di Gioia for discussions regarding LPS-response in macrophages and early 594 experimental work. The work was funded by an Allen Distinguished Investigator award through 595 The Paul G. Allen Frontiers Group to N.S., a Bits to Bytes award from MLSC to N.S., an NIGMS 596 award R01GM144967 to N.S., and a MIRA award from the NIGMS of the NIH (R35GM148218) 597 to N.S. 598

Competing interests: N.S. is a founding director and CEO of Parallel Squared Technology Insti tute, which is a nonprofit research institute.

Author contributions

602	Experimental design: J.D., T.J., and N.S.
603	Data analysis: J.D.
604	LC-MS/MS: J.D. and T.J.
605	Sample preparation: J.D., T.J., A.L., L.K., S.K., and M
606	Raising funding: N.S.
607	Supervision: N.S.
608	Initial draft: J.D., T.J., and N.S.

27

.R.

Writing: All authors approved the final manuscript.

610 **References**

609

611

612

613

614

615

616

617

618

619

620

621

622

623

624

625

626

627

628

629

630

631

632

633

634

635

636

- 1. Macosko, E. Z. *et al.* Highly parallel genome-wide expression profiling of individual cells using nanoliter droplets. *Cell* **161**, 1202–1214 (2015).
 - 2. Slavov, N. Scaling Up Single-Cell Proteomics. *Molecular & Cellular Proteomics* **21**, 100179 (2022).
- 3. Quake, S. R. A decade of molecular cell atlases.
 - 4. Symmons, O. & Raj, A. Whats Luck Got to Do with It: Single Cells, Multiple Fates, and Biological Nondeterminism. *Molecular cell* **62**, 788–802 (2016).
- 5. Slavov, N. Unpicking the proteome in single cells. *Science* **367**, 512–513 (2020).
 - 6. Weinreb, C., Wolock, S., Tusi, B. K., Socolovsky, M. & Klein, A. M. Fundamental limits on dynamic inference from single-cell snapshots. *Proceedings of the National Academy of Sciences* **115**, E2467–E2476 (2018).
 - 7. Leduc, A., Harens, H. & Slavov, N. Modeling and interpretation of single-cell proteogenomic data. *arXiv*. eprint: 2308.07465 (2023).
 - 8. Slavov, N. Learning from natural variation across the proteomes of single cells. *PLOS Biology* **20**, 1–4 (Jan. 2022).
 - 9. Kelly, R. T. Single-Cell Proteomics: Progress and Prospects. *Molecular & Cellular Proteomics* (2020).
 - 10. Peters-Clarke, T. M. *et al.* Boosting the Sensitivity of Quantitative Single-Cell Proteomics with Infrared-Tandem Mass Tags. *J. Proteome Res.* (May 2024).
- 11. Peters-Clarke, T. M., Coon, J. J. & Riley, N. M. Instrumentation at the Leading Edge of Proteomics. *Analytical Chemistry*.
- 12. Slavov, N. Driving Single Cell Proteomics Forward with Innovation. *Journal of Proteome Research* **20**, 4915–4918 (2021).
- 13. Gatto, L. *et al.* Initial recommendations for performing, benchmarking, and reporting singlecell proteomics experiments. *Nat. Methods* **20**, 375–386 (2023).
- 14. Slavov, N. Single-cell protein analysis by mass spectrometry. *Current Opinion in Chemical Biology* **60**, 1–9 (2020).
- Specht, H. *et al.* Single-cell proteomic and transcriptomic analysis of macrophage hetero geneity using SCoPE2. *Genome Biology* 22, 50. (2021) (Jan. 2021).
- 64016. Cong, Y. *et al.* Ultrasensitive single-cell proteomics workflow identifies > 1000 protein641groups per mammalian cell. *Chemical Science* (2020).

642 643 644	17.	Huffman, R. G. <i>et al.</i> Prioritized mass spectrometry increases the depth, sensitivity and data completeness of single-cell proteomics. en. <i>Nature Methods</i> 20 , 714–722. (2024) (May 2023).
645 646	18.	Thielert, M. <i>et al.</i> Robust dimethyl-based multiplex-DIA doubles single-cell proteome depth via a reference channel. <i>Mol. Syst. Biol.</i> 19 , e11503 (Sept. 2023).
647 648	19.	Leduc, A., Huffman, R. G., Cantlon, J., Khan, S. & Slavov, N. Exploring functional protein covariation across single cells using nPOP. <i>Genome Biology</i> 23 , 261. (2023) (Dec. 2022).
649	20.	Orsburn, B. C., Yuan, Y. & Bumpus, N. N. Insights into protein post-translational modification landscapes of individual human cells by trapped ion mobility time-of-flight mass spectrometry. en. <i>Nature Communications</i> 13 , 7246. (2024) (Nov. 2022).
652 . 653	21.	Papalexi, E. & Satija, R. Single-cell RNA sequencing to explore immune cell heterogeneity. en. <i>Nature Reviews Immunology</i> 18, 35–45. (2024) (Jan. 2018).
654 . 655	22.	Derks, J. <i>et al.</i> Increasing the throughput of sensitive proteomics by plexDIA. en. <i>Nat. Biotechnol.</i> (July 2022).
656 . 657	23.	Derks, J. & Slavov, N. Strategies for increasing the depth and throughput of protein analysis by plexDIA. <i>Journal of Proteome Research</i> 22 , 697–705 (2023).
658	24.	Budnik, B., Levy, E., Harmange, G. & Slavov, N. SCoPE-MS: mass spectrometry of single mammalian cells quantifies proteome heterogeneity during cell differentiation. <i>Genome Biology</i> 19 , 161. (2024) (Oct. 2018).
661 . 662 663	25.	Specht, H. & Slavov, N. Optimizing Accuracy and Depth of Protein Quantification in Experiments Using Isobaric Carriers. <i>Journal of Proteome Research</i> 20 , 880–887. (2021) (Jan. 2021).
664 665	26.	Welter, A. S. <i>et al.</i> Combining data independent acquisition with spike-in SILAC (DIA-SiS) improves proteome coverage and quantification. <i>bioRxiv</i> (2024).
666 . 667	27.	Van Bentum, M. et al. Spike-in enhanced phosphoproteomics uncovers synergistic signaling responses to MEK inhibition in colon cancer cells en. May 2024. (2024).
668 . 669	28.	Tay, S. <i>et al.</i> Single-cell NF- κ B dynamics reveal digital activation and analogue information processing. <i>Nature</i> 466 , 267–271 (2010).
670 . 671	29.	Son, M. <i>et al.</i> Spatiotemporal NF- κ B dynamics encodes the position, amplitude, and duration of local immune inputs. <i>Sci Adv</i> 8 , eabn6240 (Sept. 2022).
672 673	30.	Levy, E. & Slavov, N. Single cell protein analysis for systems biology. <i>Essays In Biochemistry</i> 62 (4 2018).
674 675	31.	Wang, A. G., Son, M., Kenna, E., Thom, N. & Tay, S. NF- κ B memory coordinates transcriptional responses to dynamic inflammatory stimuli. <i>Cell reports</i> 40 (2022).
676 677	32.	Zhang, Q. <i>et al.</i> NF-κB dynamics discriminate between TNF doses in single cells. <i>Cell systems</i> 5 , 638–645 (2017).
678 679 680	33.	Lee, R. E. C., Walker, S. R., Savery, K., Frank, D. A. & Gaudet, S. Fold change of nuclear NF-B determines TNF-induced transcription in single cells. eng. <i>Molecular Cell</i> 53 , 867–879 (Mar. 2014).

- 68134.Noursadeghi, M. *et al.* Quantitative imaging assay for NF-kappaB nuclear translocation in
primary human macrophages. eng. Journal of Immunological Methods **329**, 194–200 (Jan.
2008).
- 68435.Sakai, N. *et al.* Direct Visualization of the Translocation of the -Subspecies of Protein Kinase685C in Living Cells Using Fusion Proteins with Green Fluorescent Protein. Journal of Cell686Biology 139, 1465–1476. (2024) (Dec. 1997).

687

688

689

690

691

692

693

694

695

696

697

698

699

700

701

702

703

704

705

706

707

708

709

710

711

712

713

- Wang, Y., Shyy, J. Y.-J. & Chien, S. Fluorescence Proteins, Live-Cell Imaging, and Mechanobiology: Seeing Is Believing. en. *Annual Review of Biomedical Engineering* 10, 1–38. (2024) (Aug. 2008).
- 37. Krishnaswami, S. R. *et al.* Using single nuclei for RNA-seq to capture the transcriptome of postmortem neurons. en. *Nature Protocols* **11**, 499–524. (2024) (Mar. 2016).
- Ammar, C., Gruber, M., Csaba, G. & Zimmer, R. MS-EmpiRe Utilizes Peptide-level Noise Distributions for Ultra-sensitive Detection of Differentially Expressed Proteins[S]. English. *Molecular & Cellular Proteomics* 18, 1880–1892. (2024) (Sept. 2019).
- 39. Bartels, R. The Rank Version of von Neumann's Ratio Test for Randomness. *Journal of the American Statistical Association* **77**, 40–46. (2024) (1982).
- 40. Ghavami, A., van der Giessen, E. & Onck, P. R. Energetics of Transport through the Nuclear Pore Complex. *PLoS ONE* **11**, e0148876. (2024) (Feb. 2016).
- 41. Wing, C. E., Fung, H. Y. J. & Chook, Y. M. Karyopherin-mediated nucleocytoplasmic transport. en. *Nature Reviews Molecular Cell Biology* **23**, 307–328. (2024) (May 2022).
 - 42. Keminer, O. & Peters, R. Permeability of single nuclear pores. *Biophysical Journal* **77**, 217–228. (2024) (July 1999).
- 43. Paci, G., Zheng, T., Caria, J., Zilman, A. & Lemke, E. A. Molecular determinants of large cargo transport into the nucleus. *eLife* **9** (eds Singer, R. H. & Pfeffer, S. R.) e55963. (2024) (July 2020).
- 44. Timney, B. L. *et al.* Simple rules for passive diffusion through the nuclear pore complex. *Journal of Cell Biology* **215**, 57–76. (2023) (Oct. 2016).
- 45. Zhou, H.-R., Islam, Z. & Pestka, J. J. Kinetics of lipopolysaccharide-induced transcription factor activation/inactivation and relation to proinflammatory gene expression in the murine spleen. *Toxicology and Applied Pharmacology* **187**, 147–161. (2024) (Mar. 2003).
- 46. Ochi, T. *et al.* PAXX, a paralog of XRCC4 and XLF, interacts with Ku to promote DNA double-strand break repair. *Science (New York, N.Y.)* **347,** 185–188. (2024) (Jan. 2015).
- 47. Xing, M. *et al.* Interactome analysis identifies a new paralogue of XRCC4 in non-homologous end joining DNA repair pathway. en. *Nature Communications* **6**, 6233. (2024) (Feb. 2015).
- 48. Liu, X., Shao, Z., Jiang, W., Lee, B. J. & Zha, S. PAXX promotes KU accumulation at DNA breaks and is essential for end-joining in XLF-deficient mice. *Nature Communications* 8, 13816. (2024) (Jan. 2017).
- 49. McCloskey, A., Ibarra, A. & Hetzer, M. W. Tpr regulates the total number of nuclear pore complexes per cell nucleus. *Genes & Development* **32**, 1321–1331. (2023) (Oct. 2018).

720 721	50.	Leduc, A., Koury, L., Cantlon, J. & Slavov, N. <i>Massively parallel sample preparation for multiplexed single-cell proteomics using nPOP</i> en. Nov. 2023. (2024).
722 723	51.	Petelski, A. A. <i>et al.</i> Multiplexed single-cell proteomics using SCoPE2. <i>Nature Protocols</i> 16 , 5398–5425 (2021).
724 725 726	52.	Van Bergen, N. J. <i>et al.</i> Pathogenic variants in nucleoporin TPR (translocated promoter region, nuclear basket protein) cause severe intellectual disability in humans. <i>Human Molecular Genetics</i> 31 , 362–375. (2024) (Feb. 2022).
727 728	53.	Mosalaganti, S. <i>et al.</i> AI-based structure prediction empowers integrative structural analysis of human nuclear pores. <i>Science</i> 376 , eabm9506. (2024) (June 2022).
729 730	54.	Mathieson, T. <i>et al.</i> Systematic analysis of protein turnover in primary cells. <i>Nature Communications</i> 9 , 689. (2024) (Feb. 2018).
731 732 733	55.	Savas, J. N., Toyama, B. H., Xu, T., Yates, J. R. & Hetzer, M. W. Extremely Long-lived Nuclear Pore Proteins in the Rat Brain. <i>Science (New York, N.y.)</i> 335, 942. (2024) (Feb. 2012).
734 735 736	56.	Raman, N., Weir, E. & Müller, S. The AAA ATPase MDN1 Acts as a SUMO-Targeted Regulator in Mammalian Pre-ribosome Remodeling. English. <i>Molecular Cell</i> 64 , 607–615. (2024) (Nov. 2016).
737 738 739	57.	Schink, S., Ammar, C., Chang, YF., Zimmer, R. & Basan, M. Analysis of proteome adaptation reveals a key role of the bacterial envelope in starvation survival. <i>Molecular Systems Biology</i> 18 , e11160. (2024) (Dec. 2022).
740 741	58.	Ori, A. <i>et al.</i> Cell type-specific nuclear pores: a case in point for context-dependent stoi- chiometry of molecular machines. <i>Molecular systems biology</i> 9 , 648 (2013).
742 743 744	59.	Lund, M. E., To, J., O'Brien, B. A. & Donnelly, S. The choice of phorbol 12-myristate 13-acetate differentiation protocol influences the response of THP-1 macrophages to a pro- inflammatory stimulus. <i>Journal of Immunological Methods</i> 430 , 64–70. (2023) (Mar. 2016).
745 746	60.	Specht, H. et al. Automated sample preparation for high-throughput single-cell proteomics en. Aug. 2018. (2022).
747 748 749	61.	Dou, M. <i>et al.</i> Automated Nanoflow Two-Dimensional Reversed-Phase Liquid Chromatography System Enables In-Depth Proteome and Phosphoproteome Profiling of Nanoscale Samples. <i>Analytical Chemistry</i> 91 , 9707–9715. (2024) (Aug. 2019).
750 751	62.	Meier, F. <i>et al.</i> diaPASEF: parallel accumulation–serial fragmentation combined with data- independent acquisition. en. <i>Nature Methods</i> 17 , 1229–1236. (2024) (Dec. 2020).
752 753	63.	Demichev, V. <i>et al.</i> dia-PASEF data analysis using FragPipe and DIA-NN for deep proteomics of low sample amounts. en. <i>Nature Communications</i> 13 , 3944. (2024) (July 2022).
754 755 756	64.	Demichev, V., Messner, C. B., Vernardis, S. I., Lilley, K. S. & Ralser, M. DIA-NN: neural networks and interference correction enable deep proteome coverage in high throughput. <i>Nature methods</i> 17 , 41–44 (2020).

757 758 759	65.	Cox, J. <i>et al.</i> Accurate Proteome-wide Label-free Quantification by Delayed Normalization and Maximal Peptide Ratio Extraction, Termed MaxLFQ. <i>Molecular & Cellular Proteomics</i> : <i>MCP</i> 13 , 2513–2526. (2022) (Sept. 2014).
760 761	66.	Thul, P. J. & Lindskog, C. The human protein atlas: A spatial map of the human proteome. <i>Protein Science : A Publication of the Protein Society</i> 27 , 233–244. (2021) (Jan. 2018).
762 763	67.	Pontén, F., Jirström, K. & Uhlen, M. The Human Protein Atlas—a tool for pathology. en. <i>The Journal of Pathology</i> 216 , 387–393. (2024) (2008).
764 765 766	68.	Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Power- ful Approach to Multiple Testing. <i>Journal of the Royal Statistical Society. Series B (Method-</i> <i>ological)</i> 57 , 289–300. (2024) (1995).
767 768 769	69.	Reimand, J., Kull, M., Peterson, H., Hansen, J. & Vilo, J. g:Profiler—a web-based toolset for functional profiling of gene lists from large-scale experiments. en. <i>Nucleic Acids Research</i> 35 , W193–W200. (2024) (July 2007).
770 771 772	70.	Kolberg, L. <i>et al.</i> g:Profiler—interoperable web service for functional enrichment analysis and gene identifier mapping (2023 update). en. <i>Nucleic Acids Research</i> 51 , W207–W212. (2024) (July 2023).
773 774	71.	The UniProt Consortium. UniProt: the Universal Protein Knowledgebase in 2023. <i>Nucleic Acids Research</i> 51 , D523–D531. (2024) (Jan. 2023).
775 776	72.	Johnson, W. E., Li, C. & Rabinovic, A. Adjusting batch effects in microarray expression data using empirical Bayes methods. eng. <i>Biostatistics (Oxford, England)</i> 8 , 118–127 (Jan. 2007).
777 778 779	73.	Leek, J. T., Johnson, W. E., Parker, H. S., Jaffe, A. E. & Storey, J. D. The sva package for removing batch effects and other unwanted variation in high-throughput experiments. <i>Bioinformatics</i> 28 , 882–883. (2024) (Mar. 2012).
780 781 782	74.	Makowski, D., Ben-Shachar, M. S. & Lüdecke, D. bayestestR: Describing Effects and their Uncertainty, Existence and Significance within the Bayesian Framework. en. <i>Journal of Open Source Software</i> 4 , 1541. (2024) (Aug. 2019).
783 784	75.	Goddard, T. D. <i>et al.</i> UCSF ChimeraX: Meeting modern challenges in visualization and analysis. en. <i>Protein Science</i> 27 , 14–25. (2024) (2018).

Extended Data Figures

Single-nucleus proteomics identifies regulators of protein transport

Jason Derks,^{1,2,⊠} Tobias Jonson,¹ Andrew Leduc,¹ Saad Khan,¹ Luke Khoury,¹
 Mahmoud-Reza Rafiee,¹ Nikolai Slavov^{1,2,⊠}

¹Departments of Bioengineering, Biology, Chemistry and Chemical Biology, Single Cell Center, Northeast ern University, Boston, MA 02115, USA

⁷⁹² ²Parallel Squared Technology Institute, Watertown, MA 02472, USA

 \in Data, code & protocols: scp.slavovlab.net

786

787

794



Extended Data Fig. 1 | **Evaluating nuclear enrichment of bulk samples**. Proteins markers from The Human Protein Atlas were used to benchmark the enrichment of nuclear proteins, and the depletion of non-nuclear proteins in nuclear lysate relative to whole-cell lysate.



Extended Data Fig. 2 | Assessing the continuity of time-series data for LPS-induced differentially abundant proteins. a Calculation for the ranked version of von Neumann's ratio (RVN). b Distribution of RVN ratios for differentially abundant proteins (5% FDR) in nuclei ("Sig. proteins") compared to a simulated null model. c Same as b, but for whole-cells.



Extended Data Fig. 3 | **Fold-change of LPS-induced protein transport is mass-dependent**. Absolute value fold-changes between LPS-treated and NT nuclear proteomes, and molecular masses are plotted for differentially abundant proteins (5% FDR). Smaller proteins tend to have greater fold-changes than larger proteins. Spearman correlations and associated statistical significance are noted.



Extended Data Fig. 4 | Change in the nuclear abundance of proteins associated with DNA repair complex in response to 1μ g/mL LPS. Time series trends of DNA repair complex proteins computed from up to 6 bulk biological replicates. Proteins shown in red are differentially abundant at 5% FDR, otherwise they are shown in black.



Extended Data Fig. 5 | **Benchmarking SCoPE-DIA for single nucleus proteomics**. **a** We sought to test whether acquiring single nucleus data in parallel with a more abundant carrier sample may improve protein coverage with minimal effect to quantitative accuracy. To test this, a mixed-species spike-in of 4:1 *S. cerevisiae* and 1:1 *H. sapiens* was created at 0x, 1x, 5x, 10x, 25x, and 50x carrier amounts. **b** Quantitative accuracy of proteins quantified across all carrier-levels (n=658) remains accurate despite the potential for increased interference. Dashed lines correspond to the theoretical expectation of the spike-in ratio. **c** Median ratios of precursor abundances to the carrier were computed and plotted as a histogram. Precursors with systematically compressed ratios from the theoretical expectation of the carrier level (*e.g.* 1:50) are likely poorly quantified, and were removed from downstream analysis. **d** Quantitative accuracy for intersected proteins at 4 filtering thresholds with the 50x carrier. Filtering precursors based on observed compression relative to the carrier improves quantitative accuracy. **e** Number of protein ratios quantified precursors, the 50x carrier after various filtering thresholds. **f** After filtering to remove poorly quantified precursors, the 50x carrier still enables nearly 4-fold greater proteomic coverage. The protein-level quantitative accuracy for all proteins is shown in the bottom panel. The data indicate that the overall quantitative accuracy decreases as the carrier enables identification of otherwise unidentifiable proteins; naturally, these proteins are lowly abundant and thus poorly quantified.



Extended Data Fig. 6 | **Single nucleus quality control**. **a** Single nuclei and negative controls were filtered for Translated and Channel Q-values < 0.1, and the number of precursors quantified was counted and plotted as a histogram. Successfully prepared and acquired nuclei, which were separable from negative controls, were retained for downstream analysis (highlighted in green). **b** Single nuclei were acquired with either a 25x or 50x carrier, and the median ratio of precursor intensities from single nuclei to normalized carrier was computed for each precursor. Only precursors with <6-fold compression were retained for downstream analysis (highlighted in green). **c** Nuclear purity was assessed for each single nucleus to a reference bulk nuclear fraction, shown in the y-axis, and a 3rd degree polynomial was fit for expected protein enrichment in the x-axis. Each line denotes the 3rd degree polynomial fit for a given single nucleus, and is colored by the computed area under the curve (AUC). Single nuclei with greater AUC reflect nuclei which are generally less pure (more whole-cell like). Only nuclei with AUC < 5 were retained for down-stream analysis. **d** Similar to the previous plot, this shows the distribution of AUCs for single nuclei. Only relatively pure nuclei (AUC < 5) were retained for downstream analysis.



Extended Data Fig. 7 | **Computing the transport score of single nuclei**. Single-nucleus protein transport cannot be directly quantified in our analysis due to the lack of measurements before and after LPS stimulation for the same nucleus. Here, we derive a metric, we term 'transport score,' that serves as an approximation. Deviations of protein abundances from the NT-population of single-nuclei are calculated globally, for all proteins, for each LPS-treated nucleus. The resulting vectors are weighted according to the differential protein abundances derived from the original bulk data presented in Figure 1. The mean component of the resulting vector is the weighted deviation of a single-nucleus from the NT-population of single-nuclei; we use this metric (transport score) to infer the magnitude of protein transport.



Extended Data Fig. 8 | **Protein set enrichment analysis on protein correlations to transport score**. The vector of ordered protein correlations to transport score were analyzed for enriched protein sets (Gene-Ontology terms). The results for which are shown at 1% FDR, ordered by their relative enrichment, some of which are highlighted.

Differential proteins used to compute effect of KD on transport (slope)



Extended Data Fig. 9 | **Predictive power of correlations to transport remain similar at different filtering levels**. To assess the robustness of our findings, changes in transport were either calculated based on **a** the 25 most differentially abundant proteins in the original bulk data, or **b** all differentially abundant proteins at 1% FDR (308 proteins). The trends are similar, but with decreasing predictive power as proteins with less biological signal are included for computing the slopes.



Extended Data Fig. 10 | **Protein-set enrichment analysis of whole-cells as a result of each siRNA knockdown**. The abundances of differentially enriched Gene-Ontology terms are plotted (y-axis) for each siRNA condition (x-axis); the x-axis is ordered based on each knocked-down protein's correlation to transport score. knockdowns of potential transport suppressors are generally enriched for macrophage-like processes and nucleocytoplasmic transport compared to knockdowns of potential transport enhancers.

Supplementary Figures

Single-nucleus proteomics identifies regulators of protein transport

798	Jason Derks, 1,2,oxtimes Tobias Jonson, 1 Andrew Leduc, 1 Saad Khan, 1 Luke Khoury, 1
799	Mahmoud-Reza Rafiee, 1 Nikolai Slavov $^{1,2, oxtimes}$
800	¹ Departments of Bioengineering, Biology, Chemistry and Chemical Biology, Single Cell Center, Northeast-
801	ern University, Boston, MA 02115, USA
802	² Parallel Squared Technology Institute, Watertown, MA 02472, USA
803	\in Data, code & protocols: scp.slavovlab.net

804

795

796



siRNA mediated knockdowns, whole-cells

Supplementary Fig. 1 | Volcano plots to validate siRNA mediated knockdowns in whole-cells. Differential abundance analysis comparing knockdown to negative control (non-targeting) knockdown, shown as volcano plots from the whole-cell fractions of THP-1 derived macrophages. Plots highlight and label the knocked-down protein. All knocked-down proteins are less abundant in the knockdown condition than in the negative control and 6/16 are statistically significant.



siRNA mediated knockdowns, nuclei

Supplementary Fig. 2 | Volcano plots to validate siRNA mediated knockdowns in nuclei. Differential abundance analysis comparing knockdown to negative control (non-targeting) knockdown, shown as volcano plots from the nuclear fractions of THP-1 macrophages. Plots highlight and label the knocked-down protein. All knocked-down proteins are less abundant in the knockdown condition than in the negative control and 13/16 are statistically significant.



Supplementary Fig. 3 | **Differences in protein transport for each biological replicate.** All biological replicates for each siRNA-mediated gene-knockdown are shown. Slopes for each experimental condition are calculated with respect to the original bulk data (x-axis). The difference in slopes, which we infer as the change in global protein transport as a result of the knockdown, is annotated for each biological replicate and quantified relative to its respective matched biological replicate negative control (non-targeting) siRNA.